

# **Chapter 2**

## **Basic Concepts of Probability Theory**

---

*ENCS6161 - Probability and Stochastic  
Processes*

Concordia University



# Specifying Random Experiments


- Examples of random experiments: tossing a coin, rolling a dice, the lifetime of a harddisk.
- Sample space: the set of all possible outcomes of a random experiment.
- Sample point: an element of the sample space  $S$
- Examples:  $S = \{H, T\}$   
 $S = \{1, 2, 3, 4, 5, 6\}$   
 $S = \{t | 1 < t < 10\}$
- Event: a subset of a sample space  $A \subseteq S$   
 $A = \{H\}$   
 $A = \{2, 4, 6\}$   
 $A = \emptyset$

# The Axioms of Probability

- A probability measure is a set function  $P(\cdot)$  that satisfies the following axioms.
  1.  $P(A) \geq 0$
  2.  $P(S) = 1$
  3. If  $A \cap B = \emptyset$ , then  $P(A \cup B) = P(A) + P(B)$
  4. If  $A_1, A_2, \dots$  are events s.t.  $A_i \cap A_j = \emptyset$  for all  $i \neq j$  then  $P\left(\bigcup_{k=1}^{\infty} A_k\right) = \sum_{k=1}^{\infty} P(A_k)$
- Corollary 1:  $P(A^c) = 1 - P(A)$
- Corollary 2:  $P(A) \leq 1 \quad \forall A$
- Corollary 3:  $P(\emptyset) = 0$

# The Axioms of Probability

- Corollary 4: If  $A_1 \cdots A_n$  are mutually exclusive, i.e.  $A_i \cap A_j = \emptyset, \forall i \neq j$  then  $P(\cup_{k=1}^n A_k) = \sum_{k=1}^n P(A_k)$
- Corollary 5:  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$



Union bound:  $P(A \cup B) \leq P(A) + P(B)$

# Counting Sample Points

- Permutation:  $n$  distinct objects, how many ways can we arrange them?

$$n \cdot (n - 1) \cdot (n - 2) \cdots 2 \cdot 1 = n!$$

- Selection with order: select  $k$  objects from  $n \geq k$  objects

$$n \cdot (n - 1) \cdots (n - k + 1) = \frac{n!}{(n - k)!} = {}_n P_k$$

- Combination (selection *without* order): select  $k$  objects from  $n \geq k$  objects

$$C_k^n = \frac{n!}{(n - k)!k!} = \frac{\square \square}{k}$$

# Counting Sample Points

- Example: choose a president and a treasurer from 50 students
  1. no restrictions
  2. A will serve only if he is president
  3. B & C will serve together or not at all
  4. D & E will not serve together

# Conditional Probability

- The conditional prob. of  $A$  given  $B$  has occurred is defined by :

$$P(A|B) = \frac{P(A \cap B)}{P(B)}, \quad P(B) > 0$$

Similarly,

$$P(B|A) = \frac{P(A \cap B)}{P(A)}, \quad P(A) > 0$$

So

$$P(A \cap B) = P(B)P(A|B) = P(A)P(B|A)$$

# Bayes' Rule

- If  $B_1 \cdots B_n$  are mutually exclusive and  $\cup B_i = S$ , we call these sets a partition of  $S$ .

- Theorem of total probability

For any event  $A$ , if  $B_1, \dots, B_n$  is a partition of  $S$ ,

$$\begin{aligned} P(A) &= P(A \cap B_1) + P(A \cap B_2) + \dots + P(A \cap B_n) \\ &= P(A|B_1)P(B_1) + P(A|B_2)P(B_2) + \dots + P(A|B_n)P(B_n) \end{aligned}$$

- Bayes' Rule


Let  $B_1, \dots, B_n$  be a partition of  $S$ , and  $P(A) \geq 0$ , then

$$P(B_j | A) = \frac{P(A \cap B_j)}{P(A)} = \frac{P(A|B_j)P(B_j)}{\sum_{k=1}^n P(A|B_k)P(B_k)}$$



# Bayes' Rule

- Example: binary communication system



1. Find  $P(Y = 0)$
2. Find  $P(X = 0|Y = 0)$  and  $P(X = 1|Y = 0)$

# Bayes' Rule

- Solution: let  $A$  be event  $Y = 0$ ,  $B_0$  be  $X = 0$ ,  $B_1$  be  $X = 1$

$$\begin{aligned}P(Y = 0) &= P(A) = P(A|B_0)P(B_0) + P(A|B_1)P(B_1) \\ &= (1 - \varepsilon)P + \varepsilon(1 - P)\end{aligned}$$

$$\begin{aligned}P(X = 0|Y = 0) &= P(B_0|A) = \frac{P(A \cap B_0)}{P(A)} \\ &= \frac{P(A|B_1)P(B_1)}{P(A)} = \frac{(1 - \varepsilon)P}{(1 - \varepsilon)P + \varepsilon(1 - P)}\end{aligned}$$

Similarly:

$$P(X = 1|Y = 0) = \frac{\varepsilon(1 - P)}{(1 - \varepsilon)P + \varepsilon(1 - P)}$$

# Bayes' Rule

- If  $P(X = 0|Y = 0) > P(X = 1|Y = 0)$ , we decide  $X = 0$ , i.e., if

$$(1 - \varepsilon)P > \varepsilon(1 - P) \Rightarrow P > \varepsilon,$$

we decide  $X = 0$ .

- Similarly, if  $P < \varepsilon$ , we decide  $X = 1$ .
- A special case: if  $P = 0.5$  and  $\varepsilon < 0.5$  then

$Y = 0 \Rightarrow$  we decide  $X = 0$

$Y = 1 \Rightarrow$  we decide  $X = 1$

# Bayes' Rule

- Example: 2% of people have on type of blood disease. If a person has the disease and take a blood test, with 96% probability, the result is positive and with 4% probability, the result is negative. If a person without the disease takes a blood test, then 94% negative and 6% positive. Find

$$P(\text{have disease}|\text{positive})$$

# Bayes' Rule

- Solution: Let

$D$ : a person has the disease     $D^0$ : no disease

$B$ : blood test positive             $B^0$ : blood test negative

Then  $P(D) = 2\%$  and

$$P(B|D) = 0.96 \quad P(B^0|D) = 0.04$$

$$P(B|D^0) = 0.06 \quad P(B^0|D^0) = 0.94$$


Apply Bayes Rule:

$$P(D|B) = 0.246 = 24.6\%$$

# Bayes' Rule



- We can also use tree diagram.



$$P(D|B) = \frac{P(D \cap B)}{P(B)} = \frac{0.0192}{0.078} = 0.246$$



# Bayes' Rule

- How to find  $P(D|BB)$ ?

$$\begin{aligned}P(D|BB) &= \frac{P(D \cap BB)}{P(BB)} \\&= \frac{0.02 * 0.96 * 0.96}{0.02 * 0.96 * 0.96 + 0.98 * 0.06 * 0.06} \\&= 83.9\%\end{aligned}$$

# Independence of Events

- $A$  and  $B$  are independent iff

$$P(A \cap B) = P(A)P(B)$$

- If  $A$  and  $B$  are independent

$$\begin{aligned}P(A|B) &= \frac{P(A \cap B)}{P(B)} \\ &= \frac{P(A)P(B)}{P(B)} \\ &= P(A)\end{aligned}$$

Similarly

$$P(B|A) = P(B)$$



# Independence of Events

- Events  $A_1, A_2, \dots, A_n$  are independent if

$$P(A_{i_1} \cap A_{i_2} \cdots \cap A_{i_k}) = P(A_{i_1})P(A_{i_2}) \cdots P(A_{i_k})$$

where  $1 \leq i_1 < i_2 < \cdots < i_k \leq n$

- A special case  $P(A_1 \cap \cdots \cap A_n) = \prod_{i=1}^n P(A_i)$

- In general if events are not necessarily independent

$$P(A_1 \cap A_2 \cdots \cap A_n) = P(A_1)P(A_2|A_1)P(A_3|A_1 \cap A_2) \cdots \\ \cdots P(A_n|A_1 \cap \cdots \cap A_{n-1})$$